Colville tsunami hazards




Table of Contents

PURPOSE	3
SOURCE REFERENCE	3
WHAT IS A TSUNAMI?	3
WHERE DO TSUNAMIS COME FROM?	3
FURTHER INFORMATION	3
LOCAL SOURCE TSUNAMI FROM THE KEREPEHI FAULT	4
THE KEREPEHI FAULT	4
HOW LONG DOES IT TAKE FOR LOCAL SOURCE WAVES TO ARRIVE AT COLVILLE?	4
INUNDATION MAPS: IMPACT OF LOCAL SOURCE TSUNAMI WAVES ON COLVILLE	4
Figure 3: Otautu Bay, Colville	5
Figure 4: Whangaahi, Colville	6
Figure 5: Colville township	7
REGIONAL SOURCE TSUNAMI FROM THE TONGA-KERMADEC TRENCH	8
The Tonga-Kermadec Trench	8
How long does it take for regional source waves to arrive at Colville?	8
INUNDATION MAPS: IMPACT OF REGIONAL SOURCE TSUNAMI WAVES ON COLVILLE	8
Figure 8: Otautu Bay, Colville	9
Figure 9: Whangaahi, Colville	10
Figure 10: Colville township	11
DISTANT SOURCE TSUNAMI FROM SOUTH AMERICA	12
SOUTH AMERICAN SOURCES	12
How long does it take for distant source waves to arrive at Colville?	12
INUNDATION MAPS: IMPACT OF DISTANT SOURCE TSUNAMI WAVES ON COLVILLE	12
Figure 13: Otautu Bay, Colville	
Figure 14: Whangaahi, Colville	14
Figure 15: Colville township	15

Purpose

To summarise tsunami hazards information for the Colville community:

- Where do tsunami come from?
- How long do tsunami waves take to arrive in Colville from the various sources?
- What impacts do tsunami have on the community?

Source reference

This summary draws on information contained within the following technical report:

Borrero, J.C. (2018). *Numerical Modelling of Tsunami Inundation in the Firth of Thames*. eCoast Limited, Raglan.

The full report is available here:

http://www.waikatoregion.govt.nz/tsunamistrategy

What is a tsunami?

A tsunami is a series of water waves most commonly caused by seafloor earthquakes. Tsunami waves are different to wind-generated waves in that they are a transfer of energy, and usually travel a lot further inland than wind-generated waves

Where do tsunamis come from?

Tsunamis caused by seafloor earthquakes occur most commonly around tectonic plate boundaries, particularly around the Pacific 'Ring of Fire'. Tsunamis can also occur along undersea fault lines that lie just offshore, whether associated with a plate boundary or not.

Colville has three primary sources of tsunami:

- 'Local source' from the Kerepehi Fault
- 'Regional source' from the Tonga-Kermadec Trench just off East Cape
- 'Distant source', most commonly from large earthquakes in South America.

An overview of the tsunami sources, wave arrival times and potential inundation in Colville is provided in the following pages.

Further information

Further general information about tsunami hazards is available at: http://www.waikatoregion.govt.nz/tsunami



Work to identify tsunami hazards on the Coromandel Peninsula west coast and Firth of Thames is a joint initiative between Thames Coromandel District Council and Waikato Regional Council.



Local source tsunami from the Kerepehi Fault

The Kerepehi Fault

A large earthquake along the offshore portion of the Kerepehi Fault is thought to be capable of generating a tsunami. The size of tsunami waves and their arrival time at Colville depends largely upon the size and position of the earthquake event.

Figure 1 shows the five fault segments considered in the technical report. Of the five faults considered, a magnitude (Mw) 7.1 earthquake rupturing along 16km of segment 'D2' has the largest potential impact on the Colville community. This scenario is considered the 'maximum credible event' for Colville from the Kerepehi Fault.

How long does it take for local source waves to arrive at Colville?

Assuming a maximum credible earthquake along segment 'D2', Figure 2 shows that:

- Water levels begin to rise about 15 minutes following the earthquake, and reach a peak of around 1.6m above the existing water level around 30 minutes following the earthquake
- The first wave is the largest, and water levels continue to rise and fall rapidly for at least eight hours due to on-going wave arrivals.




Figure 1: The five Kerepehi Fault segments considered within the technical report




Figure 2: Wave arrival times and inundation levels from segment 'D2'

Inundation maps: impact of local source tsunami waves on Colville

Figures 3, 4 and 5 (overleaf) show the potential inundation from a maximum credible earthquake on segment 'D2' of the Kerepehi Fault. It is important to note that the maps:

- Assume wave arrival at Mean High Water Springs (the highest level that spring tides reach on average over a period of time)
- Only show inundation of land areas that are normally above sea level.



Figure 3: Otautu Bay, Colville
Kerepehi Fault potential inundation from a maximum credible earthquake on segment 'D2'




Figure 4: Whangaahi, Colville
Kerepehi Fault potential inundation from a maximum credible earthquake on segment 'D2'




Figure 5: Colville township

Kerepehi Fault potential inundation from a maximum credible earthquake on segment 'D2'

Regional source tsunami from the Tonga-Kermadec Trench


The Tonga-Kermadec Trench

The Tonga-Kermadec Trench is a subduction zone at a convergent tectonic plate boundary, where the Pacific Plate is being subducted underneath the Australian Plate. A large earthquake along the Tonga-Kermadec Trench to the north-east of New Zealand represents the most significant near-source tsunami threat for the Eastern Coromandel Peninsula, but will also affect the Firth of Thames, including Colville.

The technical report considers that a magnitude (Mw) 8.9 earthquake rupturing along a 450km segment of the Tonga-Kermadec Trench just off East Cape to be a 'maximum credible event' (see Figure 6). This event is similar to the Tohoku earthquake and tsunami that occurred in Japan in 2011.

How long does it take for regional source waves to arrive at Colville?

Assuming the magnitude (Mw) 8.9 earthquake described above, Figure 7 shows that:






Figure 6: The 'maximum credible event' from the Tonga-Kermadec Trench. This event is a similar magnitude and nature to the 2011 Japan tsunami

• The first and second waves rise to around 1m just prior to and after the two-hour mark, but the largest wave at around 1.3m high does not arrive until four hours following the earthquake.




Figure 7: Wave arrival times and inundation levels from the Tonga-Kermadec Trench (at MHWS)

Inundation maps: impact of regional source tsunami waves on Colville

Figures 8, 9 and 10 (overleaf) show the potential inundation from a maximum credible earthquake on the Tonga-Kermadec Trench. It is important to note that the maps:

- Assume wave arrival at Mean High Water Springs (the highest level that spring tides reach on average over a period of time)
- Only show inundation of land areas that are normally above sea level.



Figure 8: Otautu Bay, Colville
Potential inundation from a maximum credible earthquake on the Tonga-Kermadec Trench




Figure 9: Whangaahi, Colville
Potential inundation from a maximum credible earthquake on the Tonga-Kermadec Trench




Figure 10: Colville township

Potential inundation from a maximum credible earthquake on the Tonga-Kermadec Trench

Distant source tsunami from South America

South American sources

Previous studies have indicated that tsunamis produced by large earthquakes along the South American Subduction Zone have the greatest impact of all the distant tsunami sources on New Zealand.

The technical report considers three scenarios from South America (see Figure 11):

- 1. The 1960 Valdivia, Chile earthquake (magnitude ~9.2)
- 2. The 1868 Arica, Chile and Southern Peru earthquake (magnitude ~9.4)
- 3. 'FF7', a theoretical variant of the 1960 Valdivia earthquake placed in Central Peru (magnitude ~9.2).

Of the three scenarios considered, the FF7 earthquake has the most impact on Colville at Mean High Water Springs, although this *impact* is far lower than the local and regional sources.




Figure 11: Map of the 'FF 7' variant of the 1960 Chilean earthquake, placed in Central Peru

How long does it take for distant source waves to arrive at Colville?

Assuming the 'FF 7' scenario, Figure 12 shows that:

- Water levels begin to rise about 15.5 hours following the earthquake, then rise and fall rapidly (4-6 times per hour) for at least 18 hours following first wave arrival at Colville
- The initial wave is small (0.2m), and the waves rise slowly to the maximum inundation level of 0.6m above the existing water level after a further three hours.




Figure 12: Wave arrival times and inundation levels from the 'FF 7' distant source scenario

Inundation maps: impact of distant source tsunami waves on Colville

Figures 13, 14 and 15 (overleaf) show the potential inundation from a maximum credible earthquake from the Central Peru region of South America. It is important to note that the maps:

- Assume wave arrival at Mean High Water Springs (the highest level that spring tides reach on average over a period of time)
- Only show inundation of land areas that are normally above sea level.



Figure 13: Otautu Bay, Colville
Potential inundation from a maximum credible earthquake from Central Peru in South America




Figure 14: Whangaahi, Colville
Potential inundation from a maximum credible earthquake from Central Peru in South America




Figure 15: Colville township

Potential inundation from a maximum credible earthquake from Central Peru in South America