Manaia tsunami hazards




Table of Contents

PURPOSE	3
SOURCE REFERENCE	3
WHAT IS A TSUNAMI?	3
WHERE DO TSUNAMIS COME FROM?	3
FURTHER INFORMATION	3
LOCAL SOURCE TSUNAMI FROM THE KEREPEHI FAULT	4
THE KEREPEHI FAULT	4
HOW LONG DOES IT TAKE FOR LOCAL SOURCE WAVES TO ARRIVE AT MANAIA?	4
INUNDATION MAPS: IMPACT OF LOCAL SOURCE TSUNAMI WAVES ON MANAIA	4
Figure 3: Manaia Bay	5
Figure 4: Te Kouma Harbour	6
Figure 5: Kirita Bay	7
REGIONAL SOURCE TSUNAMI FROM THE TONGA-KERMADEC TRENCH	8
The Tonga-Kermadec Trench	8
How long does it take for regional source waves to arrive at Manaia?	8
INUNDATION MAPS: IMPACT OF REGIONAL SOURCE TSUNAMI WAVES ON MANAIA	8
Figure 8: Manaia Bay	9
Figure 9: Te Kouma Harbour	10
Figure 10: Kirita Bay	11
DISTANT SOURCE TSUNAMI FROM SOUTH AMERICA	12
SOUTH AMERICAN SOURCES	12
How long does it take for distant source waves to arrive at Manaia?	12
INUNDATION MAPS: IMPACT OF DISTANT SOURCE TSUNAMI WAVES ON MANAIA	12
Figure 13: Manaia Bay	
Figure 14: Te Kouma Harbour	14
Figure 15: Kirita Bay	

Purpose

To summarise tsunami hazards information for the Manaia community:

- Where do tsunami come from?
- How long do tsunami waves take to arrive in Manaia from the various sources?
- What impacts do tsunami have on the community?

Source reference

This summary draws on information contained within the following technical report:

Borrero, J.C. (2018). *Numerical Modelling of Tsunami Inundation in the Firth of Thames*. eCoast Limited, Raglan.

The full report is available here:

http://www.waikatoregion.govt.nz/tsunamistrategy

What is a tsunami?

A tsunami is a series of water waves most commonly caused by seafloor earthquakes. Tsunami waves are different to wind-generated waves in that they are a transfer of energy, and usually travel a lot further inland than wind-generated waves

Where do tsunamis come from?

Tsunamis caused by seafloor earthquakes occur most commonly around tectonic plate boundaries, particularly around the Pacific 'Ring of Fire'. Tsunamis can also occur along undersea fault lines that lie just offshore, whether associated with a plate boundary or not.

Manaia has three primary sources of tsunami:

- 'Local source' from the Kerepehi Fault
- 'Regional source' from the Tonga-Kermadec Trench just off East Cape
- 'Distant source', most commonly from large earthquakes in South America.

An overview of the tsunami sources, wave arrival times and potential inundation in Manaia is provided in the following pages.

Further information

Further general information about tsunami hazards is available at: http://www.waikatoregion.govt.nz/tsunami



Work to identify tsunami hazards on the Coromandel Peninsula west coast and Firth of Thames is a joint initiative between Thames Coromandel District Council and Waikato Regional Council.



Local source tsunami from the Kerepehi Fault

The Kerepehi Fault

A large earthquake along the offshore portion of the Kerepehi Fault is thought to be capable of generating a tsunami. The size of tsunami waves and their arrival time at Manaia depends largely upon the size and position of the earthquake event.

Figure 1 shows the five fault segments considered in the technical report. Of the five faults considered, a magnitude (Mw) 7.1 earthquake rupturing along 16km of segment 'D2' has the largest potential impact on the Manaia community. This scenario is considered the 'maximum credible event' for Manaia from the Kerepehi Fault.

How long does it take for local source waves to arrive at Manaia?

Assuming a maximum credible earthquake along segment 'D2', Figure 2 shows that:

- Water levels begin to rise almost immediately following the earthquake, and reach a peak of around 2.4m above the existing water level around 10 minutes following the earthquake
- The first wave is the largest, and water levels drop rapidly following the first wave, then continue to rise and fall at much lower levels for at least eight hours due to on-going wave arrivals.




Figure 1: The five Kerepehi Fault segments considered within the technical report




Figure 2: Wave arrival times and inundation levels from segment 'D2'

Inundation maps: impact of local source tsunami waves on Manaia

Figures 3, 4 and 5 (overleaf) show the potential inundation from a maximum credible earthquake on segment 'D2' of the Kerepehi Fault. It is important to note that the maps:

- Assume wave arrival at Mean High Water Springs (the highest level that spring tides reach on average over a period of time)
- Only show inundation of land areas that are normally above sea level.




Figure 3: Manaia Bay
Kerepehi Fault potential inundation from a maximum credible earthquake on segment 'D2'




Figure 4: Te Kouma Harbour

Kerepehi Fault potential inundation from a maximum credible earthquake on segment 'D2'



Figure 5: Kirita Bay
Kerepehi Fault potential inundation from a maximum credible earthquake on segment 'D2'

Regional source tsunami from the Tonga-Kermadec Trench


The Tonga-Kermadec Trench

The Tonga-Kermadec Trench is a subduction zone at a convergent tectonic plate boundary, where the Pacific Plate is being subducted underneath the Australian Plate. A large earthquake along the Tonga-Kermadec Trench to the north-east of New Zealand represents the most significant near-source tsunami threat for the Eastern Coromandel Peninsula, but will also affect the Firth of Thames, including Manaia.

The technical report considers that a magnitude (Mw) 8.9 earthquake rupturing along a 450km segment of the Tonga-Kermadec Trench just off East Cape to be a 'maximum credible event' (see Figure 6). This event is similar to the Tohoku earthquake and tsunami that occurred in Japan in 2011.

How long does it take for regional source waves to arrive at Manaia?

Assuming the magnitude (Mw) 8.9 earthquake described above, Figure 7 shows that:






Figure 6: The 'maximum credible event' from the Tonga-Kermadec Trench. This event is a similar magnitude and nature to the 2011 Japan tsunami

• The first wave rises to 0.35m just after the two-hour mark, the second wave is the largest at 0.6m, and the waves continue around the 0.5m level for two hours more before tapering off to around 0.25m eight hours after the earthquake.




Figure 7: Wave arrival times and inundation levels from the Tonga-Kermadec Trench (at MHWS)

Inundation maps: impact of regional source tsunami waves on Manaia

Figures 8, 9 and 10 (overleaf) show the potential inundation from a maximum credible earthquake on the Tonga-Kermadec Trench. It is important to note that the maps:

- Assume wave arrival at Mean High Water Springs (the highest level that spring tides reach on average over a period of time)
- Only show inundation of land areas that are normally above sea level.




Figure 8: Manaia Bay
Potential inundation from a maximum credible earthquake on the Tonga-Kermadec Trench




Figure 9: Te Kouma Harbour

Potential inundation from a maximum credible earthquake on the Tonga-Kermadec Trench



Figure 10: Kirita Bay
Potential inundation from a maximum credible earthquake on the Tonga-Kermadec Trench

Distant source tsunami from South America

South American sources

Previous studies have indicated that tsunamis produced by large earthquakes along the South American Subduction Zone have the greatest impact of all the distant tsunami sources on New Zealand.

The technical report considers three scenarios from South America (see Figure 11):

- 1. The 1960 Valdivia, Chile earthquake (magnitude ~9.2)
- 2. The 1868 Arica, Chile and Southern Peru earthquake (magnitude ~9.4)
- 3. 'FF7', a theoretical variant of the 1960 Valdivia earthquake placed in Central Peru (magnitude ~9.2).

Of the three scenarios considered, the FF7 earthquake has the most impact on Manaia at Mean High Water Springs, although this *impact* is far lower than the local and regional sources.




Figure 11: Map of the 'FF 7' variant of the 1960 Chilean earthquake, placed in Central Peru

How long does it take for distant source waves to arrive at Manaia?

Assuming the 'FF 7' scenario, Figure 12 shows that:

- Water levels begin to rise about 16 hours following the earthquake, then rise and fall rapidly (5-6 times per hour) for at least 14 hours following first wave arrival
- The initial wave is small (0.1m), and the waves rise and fall erratically to the maximum inundation level of 0.45m above the existing water level after a further five hours.




Figure 12: Wave arrival times and inundation levels from the 'FF 7' distant source scenario

Inundation maps: impact of distant source tsunami waves on Manaia

Figures 13, 14 and 15 (overleaf) show the potential inundation from a maximum credible earthquake from the Central Peru region of South America. It is important to note that the maps:

- Assume wave arrival at Mean High Water Springs (the highest level that spring tides reach on average over a period of time)
- Only show inundation of land areas that are normally above sea level.




Figure 13: Manaia Bay
Potential inundation from a maximum credible earthquake from Central Peru in South America



Figure 14: Te Kouma Harbour

Potential inundation from a maximum credible earthquake from Central Peru in South America



Figure 15: Kirita Bay
Potential inundation from a maximum credible earthquake from Central Peru in South America